


ILLINOIS FOOD ALLERGY EMERGENCY ACTION PLAN AND TREATMENT AUTHORIZATION



NAME: _____ D.O.B: ____/____/____

TEACHER: _____ GRADE: _____

ALLERGY TO: _____

Asthma: Yes (higher risk for a severe reaction) No

Weight: _____ lbs

ANY SEVERE SYMPTOMS AFTER SUSPECTED INGESTION:

- LUNG: Short of breath, wheeze, repetitive cough
- HEART: Pale, blue, faint, weak pulse, dizzy, confused
- THROAT: Tight, hoarse, trouble breathing/swallowing
- MOUTH: Obstructive swelling (tongue)
- SKIN: Many hives over body

Or Combination of symptoms from different body areas:

- SKIN: Hives, itchy rashes, swelling
- GUT: Vomiting, crampy pain



INJECT EPINEPHRINE IMMEDIATELY

- Call 911
- Begin monitoring (see below)
- Additional medications:
- Antihistamine
- Inhaler (bronchodilator) if asthma

Inhalers/bronchodilators and antihistamines are not to be depended upon to treat a severe reaction (anaphylaxis) → Use Epinephrine.

When in doubt, use epinephrine. Symptoms can rapidly become more severe.

MILD SYMPTOMS ONLY

- Mouth: Itchy mouth
- Skin: A few hives around mouth/face, mild itch
- Gut: Mild nausea/discomfort



GIVE ANTIHISTAMINE

- Stay with child, alert health care professionals and parent.

IF SYMPTOMS PROGRESS (see above), INJECT EPINEPHRINE

- If checked, give epinephrine for ANY symptoms if the allergen was likely eaten.
- If checked, give epinephrine before symptoms if the allergen was definitely eaten.

MEDICATIONS/DOSES

EPINEPHRINE (BRAND AND DOSE): _____

ANTIHISTAMINE (BRAND AND DOSE): _____

Other (e.g., inhaler-bronchodilator if asthma): _____

MONITORING: Stay with the child. Tell rescue squad epinephrine was given. A second dose of epinephrine can be given a few minutes or more after the first if symptoms persist or recur. For a severe reaction, consider keeping child lying on back with legs raised. Treat child even if parents cannot be reached.

- Student may self-carry epinephrine
- Student may self-administer epinephrine

CONTACTS: Call 911 Rescue squad: (____) _____

Parent/Guardian: _____ Ph: (____) _____

Name/Relationship: _____ Ph: (____) _____

Name/Relationship: _____ Ph: (____) _____

Licensed Healthcare Provider Signature: _____ (Required) Phone: _____ Date: _____

I hereby authorize the school district staff members to take whatever action in their judgment may be necessary in supplying emergency medical services consistent with this plan, including the administration of medication to my child. I understand that the Local Governmental and Governmental Employees Tort Immunity Act protects staff members from liability arising from actions consistent with this plan. I also hereby authorize the school district staff members to disclose my child's protected health information to chaperones and other non-employee volunteers at the school or at school events and field trips to the extent necessary for the protection, prevention of an allergic reaction, or emergency treatment of my child and for the implementation of this plan.

Parent/Guardian Signature: _____ Date: _____

TRAINED STAFF MEMBERS

Name: _____

Room: _____

Name: _____

Room: _____

Name: _____


Room: _____

LOCATION OF MEDICATION


- Student to carry
- Health Office/Designated Area for Medication
- Other: _____

EPIPEN Auto-Injector and EPIPEN Jr Auto-Injector Directions

- First, remove the EPIPEN Auto-Injector from the plastic carrying case
- Pull off the blue safety release cap



- Hold orange tip near outer thigh (always apply to thigh)



- Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds. Remove the EPIPEN Auto-Injector and massage the area for 10 more seconds




DEY® and the Dey logo, EpiPen®, EpiPen 2-Pak®, and EpiPen Jr 2-Pak® are registered trademarks of Dey Pharma, L.P.

Twinject® 0.3 mg and Twinject® 0.15 mg Directions



Remove caps labeled "1" and "2."

Place rounded tip against outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.



SECOND DOSE ADMINISTRATION:
If symptoms don't improve after 10 minutes, administer second dose:

Unscrew rounded tip. Pull syringe from barrel by holding blue collar at needle base.




Slide yellow collar off plunger.


Put needle into thigh through skin, push plunger down all the way, and remove.



Adrenallick™ 0.3 mg and Adrenallick™ 0.15 mg Directions



Remove GREY caps labeled "1" and "2."



Place RED rounded tip against outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.

A food allergy response kit should contain at least two doses of epinephrine, other medications as noted by the student's physician, and a copy of this Food Allergy Action Plan.

A kit must accompany the student if he/she is off school grounds (i.e., field trip).